

# Multiwavelength observations of Blazars

Stefan J. Wagner <sup>1</sup>

<sup>1</sup> Landessternwarte Heidelberg  
*E-mail(SW): s.wagner@lsw.uni-heidelberg.de*

## ABSTRACT

Blazars exhibit very broad spectral energy distributions, extending over up to 20 orders of magnitude in photon energy. Blazars also vary on a wide range of time scales with power density spectra that have been measured over as many as 9 orders of magnitude in time for the best studied objects. Given these characteristics, coordinated multiwavelength observations are required to understand the physical processes in Blazars. Apart from detailed studies of a very small number of prominent targets, statistical investigations of homogenous observations are important, but difficult to assemble due to technical constraints. The current status of multiwavelength investigations will be reviewed with a special emphasis on future opportunities.

## Multiwavelength observations of Blazars

Stefan J. Wagner  
Landessternwarte  
Heidelberg  
Germany

acknowledgements D. Emmanoulopoulos, E. Ferrero, G. Pedalletti,  
O. Kurtanidze (Abastumani), G. Bicknell (ANU Canberra)  
HESS collaboration

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1

## Multiwavelength observations

Broadband studies of Blazars are monitoring studies: 3 reasons  
Scale-free spatial and temporal structures  
Localizing emitting regions  
Clues on Lorentz factors, Uncertainties in Doppler corrections



Special emphasis:  
"Astrophysics with All-Sky X-Ray Observations"


All-Sky >> many sources >> populations (parameter studies)  
>> alerts for pointed studies  
>> long-term studies >> flux/sensitivity is 'only' bias  
(window function well understood)  
>> avoid coincidences

From examples to samples: avoiding biases, enabling parameter studies

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 2


## SED studies and timing: 1<sup>st</sup> reason



3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 3


## SED studies and timing: 1<sup>st</sup> reason



3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 4


## SED studies and timing: 2<sup>nd</sup> reason



3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 5


## SED studies and timing: 2<sup>nd</sup> reason



3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 6

## SED studies and timing: 3<sup>rd</sup> reason



Aharonian et al. (HESS collaboration, 2006)

simultaneous (single epoch) SEDs do not provide a unique constraint for models of acceleration and radiative processes.  
(It has not been demonstrated, but there is hope that time-resolved multifrequency studies will provide unique constraints.)

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 7

## General motivation

Seeking correlations to identify sources  
(when known coincidences are insufficient)


1960s finding radio sources: "3C273B = ..."  
1970s finding X-ray sources: "Ariel ... = PKS 2155-304"  
1990 finding Gamma-ray sources: 3C66A, PKS 1406-076  
2000s finding TeV sources (see Elina Lindfors)

Finding sources in the sky;  
localizing subvolumes in the sources (in space and time);  
localizing subvolumes in parameter space ( $B$ ,  $n$ ,  $\gamma$ );  
identify and understand processes (qualitative relations between bands)

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 8

## Review:



Within synchrotron band: Radio-Optical Correlations  
Lunar Occultations (e.g. Dent)  
Flares (e.g. van der Laan)  
Outburst and VLBI components (Babadzhanyants)  
Polarisation characteristics (Kikuchi)  
but: IDV (SW and Quirrenbach)  
relation to X-ray bands (e.g. Urry; Brinkmann; Courvoisier; Edelson)

Within HE bump: 3C279 (Wehrle)


Synchrotron – HE bump: several in 1980s (without mentioning)  
involving gamma-rays (1406-076; Mkn 421; Takahashi & SW)

but several counter-examples ("orphan" events); poor coverage in energy

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 9

## Parameter Space




Originally shown at 1<sup>st</sup> MAXI workshop, Nikko, 2001

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1

## Long, continuous light-curves



uninterrupted, unbiased, but coarse binning, few sources


D. Emmanoulopoulos & SW (2008)

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1

## Self-similar structures?

Mrk 421, XTE (2-10 keV)



factors of 30: 3000d to 3h

D. Emmanoulopoulos & SW, 2008

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1


## Spatial and temporal scales

Temporal scales:  
Mrk 421, X-rays: 100 s - 10 Msec  
3C345, optical: 300 s - 1 Gsec

Spatial scales:  
M87, radio (GHz): 10 mpc - 100 kpc

self-similar structures

self-similar flares  
(power-law PDS)



3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1


## Small scales

Variability as a tool to small spatial scales  
Fast flares extend resolution to very compact regions.  
Strongly structured jets  
(Single blobs in VLBI resulting from limited dynamic range).

Mrk 501 shows many small clouds.

If the self-similarity extends to smaller spatial scales, small clouds might refer to fast flares.

Extreme examples from IDV and VHE.




Giovannini et al., 2000

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1


## Small scales: Intraday Variability

Intraday variability (IDV)  
S5 0716+714 ( $z \sim 0.3$ )  
Radio (5, 8.4 GHz) and optical (650 nm)  
variations on timescales  $\sim 100$  ksec



Wagner et al., 1994

implied linear dimensions:  
500 times smaller than (resolved) vlbi knots,  
100 times smaller than unresolved vlbi core  
(assumed to be self-absorbed).  
Variable subvolume one million times smaller  
than any resolved structures  
(radiation density 100 000 times larger)



Krichbaum et al., 2001

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1


## Just how small can they be?



Emitting volume:  $R\mathcal{D} < 4.6E12$  cm

Cross-section of emitting volume:  $\pi^2 E-11$  A (A= cross-section of jet)

[Inferences in both cases:  $\mathcal{D}$  might be as high as 100]



Aharonian et al. (H.E.S.S. collaboration, 2007)

PKS 2155-304 (flare in 2006)  
time-scale 60 sec ... 300 sec  
linear scale from causality

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1

## Filling factors and duty cycles

Subvolumes producing 0.1 to 10 times the quiescent flux  
with individual filling factors  $E-10...E-15$  should result in large  
numbers of such subvolumes blending the integrated light curves.

The contrast to observations implies one of three possibilities:

- 1) Duty cycle is very low
- 2) Zone of variations extends over small radial range only
- 3) Distribution function in some crucial parameter very steep  
(highly nonlinear system)




(1) is essential a special case of (3)

consider (2): Where do Blazars produce their emission?


3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 1


## Emission site in M 87



Aharonian et al. (HESS), 2006




3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008



Harris et al., 2005




Cheung et al., 2007



Stefan Wagner: Multiwavelength Observations of Blazars 1

## Locating site of PKS 2155-304

**Evidence from spectral monitoring: Non-detection of spectral changes during bright VHE flare (Costantini et al., 2008) implies minimum distance  $> 100 R_g$**



Bicknell &amp; SW, 2008




Stefan Wagner: Multiwavelength Observations of Blazars 1

simple worries  $\gamma_{\min}, \gamma_{\max}, \Gamma$

Almost all SED “fits” of Blazars invoke high  $\gamma$  min.  
 There are other observations/problems that suggest that this is required.  
 (e.g. Blundell et al., 2004, Tsang & Kirk, 2007, SW, 1997)  
 There are very few (no?) acceleration mechanisms achieving this.



Optical and Xray observations can derive those:




Optical/XMM studies of 0716+714 (Ferrero et al., 2006)

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars

simple worries  $\gamma_{\min}, \gamma_{\max}, \Gamma$

MeV Blazars (Bloemen et al., 1995), GeV Blazars, TeV Blazars  
E(max) in Synchrotron/IC changes by  $<3>7$  orders of magnitude



PKS 0522-611/PKS 0506-612 vs. 1ES1101-232 (Aharonian et al., HESS, 2007)

simple worries  $\gamma_{\min}, \gamma_{\max}, \Gamma$ 

observed SED needs to be corrected for various excesses (host galaxies, extended emission), various absorptions (SSA, E, n\_H, pair absorption), redshift, and ... Doppler factor(s).

Notoriously difficult: VLBI vs. variability (IDV, VHE)  $\mathcal{D} \sim 1 \dots 100$   
 large values possible in exceptional sources (IDV of various kinds)  
 but difficult for large classes (TeV Blazars). [Remember M87]  
 ... if TeV Blazars they are: PKS 2155-304 200 GeV - ( $\mathcal{D} \sim 100$ )  $> 2$  GeV

Radial or lateral stratification, divergent trajectories?  
Different parts of the jets (different subvolumes, flares, wavelengths)  
may be subject to different Doppler corrections.

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars

## Summary

Simultaneous multifrequency monitoring required to compare oranges (or the right season) to oranges and (hopefully) break the degeneracy of different acceleration/radiation mechanisms.

Variability across the EM spectrum occurs on a wide range of timescales.

Individual components/flares relate to regions of very different sizes (but they are not very different otherwise).

Very small filling factors of emissivity. Stratified media.  
Single-zone models ought to be self-consistent

Multifrequency observations may also provide localisations

Range of Lorentz factors may be constrained with Maxi observations.

Doppler factors might be diverse and provide the biggest challenge.

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars 2

*The End*

Thank you for your attention

3<sup>rd</sup> Intl. Maxi Workshop, RIKEN, June 2008

Stefan Wagner: Multiwavelength Observations of Blazars